

- Generally speaking, what is the difference between the terms *intramolecular* and *intermolecular* forces of attraction?
- Is a covalent bond considered to be an *intramolecular* or an *intermolecular* force of attraction?
- What is the difference between a:
 - Non-polar covalent bond and a polar covalent bond?
 - Non-polar covalent molecule and a polar covalent molecule?
- The length of a covalent bond between two identical atoms is equivalent to the sum of their atomic radii. (*Atomic radius is the distance from the nucleus to the outermost electron.*)

For example, the atomic radius for hydrogen is 37 pm and that for fluorine is 64 pm.

Based on the above information, what would be the length of the bond for H₂, a non-polar covalent molecule?

Based on the above information, what would be the length of the bond for F₂, another non-polar covalent molecule?

What then, would be the theoretical length for the H-F bond?

Experimental evidence shows that the bond length for HF is 92 pm. What gives? Hmmm....
HF is a very polar covalent bond. Based on this fact alone, how might we explain the difference between the theoretical and the actual bond lengths? That is, what would make the bond length shorter than predicted?

- There are three types of intermolecular forces of attraction: *dispersion*, *dipole-dipole*, and *hydrogen bonding*. What are the differences between these three?
- For each of the following substances:
 - determine if the bonds are polar
 - state and sketch the shape of the molecule
 - indicate whether or not the molecule is polar
 - indicate the type of intermolecular force that would predominate (rule)

F ₂	SCl ₂	CH ₂ O	NO ₃ ¹⁻	PCl ₄ F	Br ₂	PH ₃	NBr ₃
CCl ₄	H ₂ Se	NO ₂ ¹⁻	BrF ₃	BrF ₅	SCl ₄	SnCl ₅ ¹⁻	NH ₄ ⁺

- Molecular nitrogen, N₂, has a boiling point of -196°C while oxygen, O₂, has a boiling point of -183°C.
 - Identify each molecule as being polar or non-polar.
 - Are the intermolecular forces of attraction dispersion, dipole, or hydrogen bonding?
 - Account for the differences in their boiling points.
- The boiling point for nitrogen monoxide is -157°C.
 - Is this molecule polar or non-polar?
 - Account for the fact that its boiling point is higher than those of its respective elements.
- Which would have the higher boiling point: chlorine gas or iodine monochloride? Explain.

10. Dry ice, CO_2 , is held together by what type of intermolecular forces? How does this account for the fact that dry ice readily changes from a solid to a gas?
11. Vapour pressure is the pressure that vapours would exert in a container. A substance with a high vapour pressure would evaporate more readily than one with a low vapour pressure. Which of the following substances would you expect to have the lowest vapour pressure: NH_3 , PH_3 , CH_4 ?
12. Predict which of the following substances in each pair will have the highest melting point?
- | | |
|-------------------------------------|--|
| a. CS_2 or CCl_4 | f. HI or KI |
| b. Cl_2 or F_2 | g. Na_2O or H_2O |
| c. SiO_2 or CO_2 | h. CH_4 or NH_3 |
| d. CHCl_3 or CF_4 | i. CaF_2 or HF |
| j. BF_3 or P_4 | |
13. Rank each of the following from strongest to weakest intermolecular forces:
- | |
|--|
| a. He , NH_3 , NF_3 , NaCl |
| b. HF , F_2 , FCl |
14. Identify and explain which molecule (from each pair) would have the highest melting point.
- | |
|--|
| a. CS_2 vs CCl_4 |
| b. HI vs HBr |
| c. Cl_2 vs F_2 |
| d. Na_2O vs H_2O |
| e. CH_4 vs NH_3 |
| f. CHCl_3 vs CF_4 |